Table I. Cumulative Cancer Risk to Age 70 in Individuals with Lynch Syndrome

Cancer	MLH1	MSH2	MSH6	PMS2
Colon	41-58%	48-51%	10-22%	15-20%
Endometrium	29-54%	21-24%	8-32%	15%
Ovary	6-20%	8-24%	1-11%	**
Stomach	6-11%	0.2-8%	<u>≤</u> 3%	**
Urinary tract	0.2-3%	2-4%	<1%	**
Small bowel	0.4-5%	1.1-1.3%	Not reported	**
Brain/CNS	0.3-2%	3-6%	Not reported	**
Hepatobiliary tract	2-3%	0.02-0.4%	Not reported	**
Sebaceous neoplasms	1-9%*	1-9%*	Not reported	Not reported
Pancreas	1-6%*	1-6%*	Not reported	Not reported

^{*}Risk estimate includes patients with *MLH1* and/or *MLH2* mutations

Barrow E, Robinson L, Alduai, W, et al. Cumulative lifetime incidence of extracolonic cancers in Lynch syndrome: a report of 121 families with proven mutations. Clinical Genetics 75:141-149, 2009.

Barrow E, Alduaij W, Robinson L, et al. Colorectal cancer in HNPCC: cumulative lifetime incidence, survival and tumour distribution. A report of 121 families with proven mutations. Clinical Genetics 74:233-242, 2008.

Bonadona V, Bonaiti B, Olschwang S, et al: Cancer risks associated with germline mutations in MLH1, MSH2, and MSH6 genes in Lynch syndrome. JAMA 305:2304-2310, 2011.

National Comprehensive Cancer Network: NCCN Practice Guidelines in Oncology: Genetic/Familial High-Risk Assessment: Colorectal V.1.2014. Available at www.nccn.org

Senter L, Clendenning M, Sotamaa K, et al: The clinical phenotype of Lynch syndrome due to germ-line PMS2 mutations. Gastroenterology 135:419-428, 2008.

Watson P, Vasen HF, Meckli, J, et al. The risk of extra-colonic, extra-endometrial cancer in the Lynch syndrome. International Journal of Cancer 123:444-449, 2008.

^{**6%} combined cumulative risk for renal pelvis, stomach, ovary, small bowel, ureter, brain